


Penile length augmentation

- 🕒 **Duration: 1h**
- 🪚 **Anesthesia : local with sedation**
- 🏥 **Hospitalization : no**
- 👁️ **Results : immediate**
- 📅 **Recovery : 7-10 days**
- 💰 **From 4 400 CHF**

The length of the penis can be increased by 2 to 4 cm at rest, by cutting the anterior part of the suspensory ligament. This increase in length will only be seen to a very limited extent during erection, unless liposuction of the pubic area is associated with it.

It is also possible to increase the circumference of the penis, thanks to hyaluronic acid or fat injections, by 2 to 3.5 cm. This difference will be seen significantly at rest and more discreetly when erect.



Technique :

Local anesthesia. Incision of 3 cm in the shape of an inverted V at the base of the penis, hidden in the hairy area and in the natural shadows. The anterior part of the suspensory ligament is freed and incised. Closure, then placement of biological surgical glue.

Sometimes, liposuction of the pubic area is associated to optimize the result.

Results :

Time to final results: 4-7 days (hematoma, swelling)

Longevity of results : definitive

Before surgery:

Stop smoking 1 month before

Do not take Aspirin 10 days before surgery, inform Dr. Lucas of any anticoagulant medication (Sintrom, Coumadin, Eliquis, etc..) to arrange for its discontinuation or replacement.

Take Arnica capsules 5 times a day for 5 days before surgery to prevent hematoma

Shave pubic area 3 days before surgery.

Intervention :

Pain: the pain during the intervention will be almost non-existent, with an adapted anesthesia

Duration of the procedure: 60 minutes

Anesthesia: local with sedation or general if the patient wishes

Location: operating room

Ambulatory (without hospitalization)

After surgery:

Post-operative pain: 1/10 often responding to Dafalgan

Social eviction:

Swelling and hematoma for 4-7 days

Suture removal: no sutures

Return to work: immediate

Return to sport: 2-3 weeks

Instructions:**Hyaluronic acid**

Dr Lucas check-up 2 days, 2 weeks, 3 months

No sexual activity for 2 weeks

Fat

Desinfection of the scar with Merfen for 7 days twice a day.

No wearing of tight underwear and no sexual activity for 3 weeks.

Shower from the 2nd day: for 10 days, do not rub too hard, dry with a clean cloth by dabbing.

Preventive antibiotic therapy for 3 days with Augmentin 1 g twice a day.

Control Dr Lucas 2 days, 1 week, 3 weeks, 3 months.

Complications :

- Infection, hematoma
- Decreased sensitivity on the back of the penis: it comes back the first weeks
- Penile augmentation and erection: there is no consequence on the erection, following this type of intervention.

Alternatives and operations that can be combined :

We can increase the circumference of the penis, thanks to hyaluronic acid or fat injections, from 2 to 3.5 cm. This difference will be seen significantly at rest and more discreetly when erect.

Dr Dominique Lucas

Chirurgien plasticien et esthétique FMH, EBOPRAS

Ancien Chef de Clinique universitaire CHUV

Membre FMH (Fédération médicale helvétique), KCCS (Korean Society of Cosmetic Surgery), ISAPS (International Society of Aesthetic Surgery)

Drlucasplasticsurgery@gmail.com

+41 77 992 95 97

Boulevard Helvétique 26, 1207 Genève

